Progress Report on Black-headed and Common Gull Colour-ringing Study, Lough Mask, County Mayo

14th September, 2010



Mìchèal Casey

Eoin Mc Greal, NPWS, Aghinish, Ballinrobe, Co. Mayo, Ireland 094 95 42775 eoin.mcgreal@environ.ie

Introduction

Between 2006 and 2010 a colour-ringing study of both Common and Black-headed Gulls was carried out on Lough Mask, Co. Mayo, co-ordinated by the National Parks & Wildlife Service in conjunction with BTO ringing trainer, Chris Benson.

Lough Mask is designated as a candidate Special Protection Area (SPA 004062) for wild birds under EU Directive 79/409 and part of the site's qualifying interest is for breeding gulls. Both Common and Black-headed Gulls are listed as species of conservation concern due to their unfavourable conservation status in Europe and because their national breeding populations have declined by 25-50% in the last 25 years. Recent studies have added the Blacked Gull in Ireland to the red list as its breeding range may have contracted by 95% between the periods 1968-72 and 1998-2002 and that its breeding population may have fallen by some 70% over the same period. The corresponding declines for Common Gull are also of concern. Lough Mask holds a significant percentage of the national breeding population for both species.

The initial aims of the study were to establish information on longevity, dispersal patterns, site fidelity, survival rates, etc. and to act as a mechanism for monitoring overall population trends. The colour-ringing component of the study ceases in 2010 and now the main focus will be to closely monitor fluctuations in breeding numbers, productivity, depredation, time budgets and the recruitment of colour-ringed birds back to the breeding colonies. The study also functions as part of the monitoring mechanism of the overall SPA designation.

Methods

The study was confined to the ringing of pulli and was largely carried out on 2 separate colonies; Rams Island (M091599) for Common Gulls and an islet off Aghinish (M135665) for Black-headed Gulls. All ringed pulli were fitted with an engraved plastic ring in an attempt to improve the likelihood of a bird being reported. Each ring was engraved with a unique 4-digit code and each bird can therefore be identified in the field as an individual. These coded yellow rings have been fitted on the left tarsus with a corresponding BTO metal ring attached to the right tarsus.

The following colour-ring codes were used in the study;

2006:	201A-200A . 201C-200C . 201E-200E . 201X-237X, 241X-250X, 261X-200X	Black-headed Gull Black-headed Gull Black-headed Gull Black-headed Gull	100 100 100 <u>87</u> 387
	201Z-282Z	Common Gull	82
2007:	238X-240X, 251X-260X 201H-200H 201M-200M 201P-200P 201S-200S	Black-headed Gull Black-headed Gull Black-headed Gull Black-headed Gull Black-headed Gull	13 100 100 100 <u>100</u> 413
	283Z-290Z 201V-200V	Common Gull Common Gull	8 100
2008:	201B – 200B 201D – 200D 201F – 271F	Black-headed Gull Black-headed Gull Black-headed Gull	108 100 100 <u>71</u> 271
	201Y – 271Y	Common Gull	71
2009:	272F – 200F 201G – 200G	Black-headed Gull Black-headed Gull	29 100 129
	272Y – 288Y	Common Gull	17
2010:	289Y – 200Y 291Z – 200Z	Common Gull Common Gull	12
Total:		Black-headed Gull Common Gull	1200 300

Discussion:

To date (14/09/10) 540 re-sightings of 176 post-fledged gulls have been submitted – these were made up of 359 re-sightings of 113 individual Black-headed Gulls and 181 resightings of 63 individual Common Gulls.

Away from the breeding colonies all 137 re-sightings of 48 Common Gulls received to date have been coastal and have occurred between Killala Bay and Tralee Bay, within160km of the breeding site, in a mainly west to south direction. This represents 16.7% of all Common Gull pulli ringed to 2009. To date it appears that dispersal is heavily concentrated along the South Galway and Clare coastlines. Although this colony is near the southern extremity of its breeding range the lack of any re-sightings north of County Mayo may indicate that Common Gulls breeding in western Ireland are separate from the metapopulation based in western Scotland.

331 re-sightings of 95 Black-headed Gulls away from the breeding colony have occurred over a broader front and at greater distances (7 individuals have been re-sighted in Spain up to 1650km from their natal colony) than Common Gulls but still indicate a mainly southerly dispersal pattern. This represents 7.9% of all Black-headed Gull pulli ringed to 2009 and is considerably lower than that received for Common Gulls. This may imply that a significant proportion of Black-headed Gulls are dispersed in areas that are at present under recorded or that their behaviour or habitat preferences make them more difficult to scope. It may also indicate greater mortality during the post-fledging period.

Both Black-headed and Common Gulls are known to be generally faithful to their natal site and do not mature until their 2nd or 3rd year respectively. To date 25 Common Gulls ringed as pulli have been recorded as adults during the breeding season on Lough Mask. This represents 8.7% of pulli ringed until 2009. Over the same period 23 adult Black-headed Gulls ringed as pulli were recorded back at their natal colony that represents only 1.9% of pulli ringed until 2009. The disparity in these figures may indicate that Common Gulls are experiencing higher post-fledging survival rates or that they are displaying a greater fidelity towards their natal site. The high vegetation and low outline of the Black-headed Gull colony also makes observance of colour-ringed birds much more difficult at this site. The observance of two Black-headed Gulls at the breeding colony that were previously re-sighted in NW Spain indicates that a proportion of this population are capable of dispersing long distances from their natal colony before returning as breeding adults.

The colour-ringing component of the project was concluded during the 2010 breeding season and it is hoped that further re-sightings of individually marked birds over the coming years will allow sufficient data to be collated in a final report. Initial findings indicate that post-fledging dispersal for both species occurs largely in western and southern Ireland but that a small proportion of Black-headed Gulls disperse much greater distances. Individuals from both species have shown strong fidelity to wintering sites in consecutive years. Some adult Common Gulls have returned to breeding sites on Lough Mask other than their natal colonies. The apparent disparity in survival rates between the two species is not yet readily explainable.

Acknowledgments:

Special thanks are extended to everybody listed below who has contributed to this project since its commencement in 2006, in particular to NPWS regional management who have provided both staff and financial resources, to those who have helped with the actual ringing and, especially, to those who have submitted re-sighting records;

Alekandra Borawska, Mark Breaks, Dermot Breen, Kane Bride, Juan Cabeza, Sue Callaghan, Padraig Canny, Mîcheàl Casey, Tom Chadwick, Kathie Claydon, Antonio Fernandez Cordeiro, Micheal Cowming, Pierre Crouzier Tom Cuffe Michael Davis, Joseph Devlin, Michael Doolan, Martin Enright, Sèamus Feeney, Manuel Figueiras, Seàn Geraty, Carlos Gutierrez, Iain Hill, Rob Holloway, Geoff Hunt, Jackie Hunt, Harry Hussey, Aiden Kelly, James Kilroy, Bernard King, James Knight, Keith Langdon, Ruth-Ann Leak, Larry Leneghan, Gosia Lipska, Pat Lonergan, Nils Helge Lorentzen, Gabriel Martin, Grahame McElwaine, Stephen McEvoy, Derek McLoughlin, Dave McNamara, Richard Mills, Dominic Mitchell, Cameron Moore, Jorge Mourinho, Rick Mundy, John Murphy, Irene O'Brien, Michael O'Connell, Aonghùs Ò'Donaill, Ger O'Donnell, Art Ò'Suìllebhàn, KT Pedersen, Chris Peppiat, Brendan Power, Ken Preston, Dirk Raes, Nicola Reddy, Encama Gonzalez Rodino, Juan M. Ruiz, Tono Salazar, LJ Salaverri Leiras, Robin Sandham, John Small, Moray Souter, Jon Stace, Wendy Stringer, Melinda Swann, Piotr Tadeusz, Simon Taylor, Marianne Ten Cate, Tom Tarpey, Paul Troake, Alyn Walsh, Paul Walsh, Dermot Ward, Neil Warnock, Todd Watkins, John Wells.

All further re-sightings will be gratefully received and may be submitted to either Eoin Mc Greal (eoin.mcgreal@environ.ie) or Chris Benson (illaune@eircom.net).

Selected Recaptures

Black-headed Gull

Ring Number		Age/ Type		Date	Init	Details
EL75617	Ν	1		31/05/06	EM	lough mask (unnamed) island, Mayo
(217C)	S	5	Sighted	12/01/09	MS	Bray Harbour, Wicklow
	S	5	Sighted	03/07/10	SME	Kilcoole, Co. Wicklow
EL75556	N	1		31/05/06	DMC	lough mask (unnamed) island, Mayo
(256A)	S	5	Sighted (R)	07/02/07	MC	Sligo Town (90 km, NE, 252 days)
	S	5	Sighted	08/04/07	MD	Galway Docks (44 km, SSE, 312 days)
	S	5	Sighted	21/08/07	MD	Galway Docks (44 km, SSE, 1 yr 82days)

EL75795 (255X)	N S S S S S S S	1 3 3 5 5 2 2 2	Sighted Sighted Sighted Sighted Sighted Sighted Sighted	04/06/07 12/10/07 25/11/07 01/12/07 20/02/08 30/03/08 28/05/09 26/01/10 08/03/10	EM RM RM RM MK HH EM KP	lough mask (unnamed) island, Mayo The Lough, Cork (201 km, SSE, 130 days) The Lough, Cork (201 km, SSE, 174 days) The Lough, Cork (201 km, SSE, 180 days) The Lough, Cork (201 km, SSE, 261 days) The Lough, Cork (201 km, SSE, 290 days lough mask (unnamed) island, Mayo The Lough, Cork The Lough, Cork
EW25518 (218H)	N R S	1 1 5	Recaptured Sighted	04/06/07 10/06/07 03/05/08	EM IO EGR	lough mask (unnamed) island, Mayo lough mask (unnamed) island), Pontevedra, Spain (1245 km, S, 334 days)
	S	2	Sighted	28/05/09	EM	lough mask (unnamed) island, Mayo
EW25630 (230M)	N S S S	1 2 2 2 2		10/06/07 30/05/09 02/06/09 10/01/10 21/03/10	EM EM EM MF MF	lough mask (unnamed) island, Mayo lough mask (unnamed) island, Mayo lough mask (unnamed) island, Mayo Quilmas, Galicia, Spain Quilmas, Galicia, Spain
EW25730 (230P)	N S S S S S	1 2 2 2 2 2 2 2	Sighted (R) Sighted (R) Sighted (R) Sighted (R) Sighted (R) Sighted (R)	16/06/07 19/11/07 01/12/07 20/04/08 06/10/08 10/11/08 24/01/09	EM AC AC AC JM JOM	lough mask (unnamed) island, Mayo Pontevedra, Spain (1263 km, S, 156 days) Pontevedra, Spain (1263 km, S, 168 days) Pontevedra, Spain (1263 km, S, 308 days) Ennis, Co. Clare (98 km, S, 473 days) Pontevedra, Spain (1263 km, S, 508 days) Pontevedra, Spain (1263 km, S, 584 days)
EW25832 (232S)	N S S S S	1 2 2 2 2 2	Sighted Sighted Sighted Sighted Sighted	21/06/07 30/12/08 05/01/09 18/01/09 08/02/09 02/06/09	EM LJ CG LJ LJ EM	lough mask (unnamed) island, Mayo Lugo, Spain (1200 km, S, 568 days) Lugo, Spain (1200 km, S, 573 days) Lugo, Spain (1200 km, S, 586 days) Lugo, Spain (1200 km, S, 606 days) lough mask (unnamed) island, Mayo
EW25844	N	1		21/06/07	EM	lough mask (unnamed) island, Mayo
(244S)	S	5	Sighted	26/03/09	AW	North Slob, Wexford (238 km, SE, 272 days)
	S	2	Sighted	02/06/09	EM	lough mask (unnamed) island, Mayo
	S	2	Sighted	28/06/10	EM	lough mask (unnamed) island, Mayo
EW67052 (252B)	N S	1 2	Sighted	06/06/08 05/01/10	EM ST	lough mask (unnamed) island, Mayo Swanpool Lake, Falmouth, Cornwall

Common Gull

EL75718	N	1		03/06/06	EM	Rams Island, Mayo
(218Z)	S	3	Sighted	15/10/07	MD	Galway Docks (38 km, SSE, 72 days)
, ,	S	2	Sighted	30/04/09	EM	Rams Island, Mayo
			8			
EL75732	N	1	03/06/06	EM	Rams 1	Island, Mayo
(232Z)	S	2	Sighted	26/12/07	MD	Quilty, Clare (83 km, S, 1 yr 206days)
	S	2	Sighted	30/04/09	EM	Long Rock, L. Mask, Mayo
						·
EL75752	N	1		14/06/06	EM	Rams Island, Mayo
(252 Z)	S	3	Sighted	28/07/06	MD	Galway Docks (38 km, SSE, 44 days)
, ,	S	3	Sighted	30/07/06	MD	Waterside, Galway (38 km, SSE, 46 days)
	S	3	Sighted	01/08/06	MD	Waterside, Galway (38 km, SSE, 48 days)
	S	3	Sighted	11/08/06	MD	Galway Docks (38 km, SSE, 58 days)
	S	3	Sighted	18/08/06	MD	Galway Docks (38 km, SSE, 65 days)
	S	3	Sighted	04/09/06	MD	Galway Docks (38 km, SSE, 82 days)
	S	3	Sighted	09/09/06	MD	Waterside, Galway (38 km, SSE, 87 days)
	S	3	Sighted	08/11/06	MD	Galway Docks (38 km, SSE, 147 days)
	S	5	Sighted	23/01/07	MD	Waterside, Galway (38 km, SSE, 223 days)
	S	2	Sighted	21/04/07	MD	Galway Docks (38 km, SSE, 311 days)
	S	2	Sighted	16/07/07	TC	Galway Docks (38 km, SSE, 1 yr 32days)
	S	2	Sighted	28/09/07	EM	Waterside, Galway (38 km, SSE, 1 yr 106days)
	S	2	Sighted	10/10/07	EM	Galway Docks (38 km, SSE, 1 yr 118days)
	S	2	Sighted	28/02/08	MD	Waterside, Galway (38 km, SSE, 1 yr 259days)
	S	2	Sighted	31/07/08	MD	Waterside, Galway (38 km, SSE,)
	S	2	Sighted	30/05/09	EM	Rams Island, Mayo
EW26000	N	1		04/06/07	EM	Rams Island, Mayo
(200V)	S	2	Sighted	05/08/09	DB	Bunlough Strand, Louisburgh, SW Mayo
(2001)	S	2	Sighted	30/05/09	EM	Rams Island, Mayo
	S	2	Sighted	22/07/09	EM	Bunowen, SW Mayo
	5	_	Signica	22/01/00	Livi	Ballowell, 5 W May o
EW25955	N	1		16/06/07	EM	Rams Island, Mayo
(255V)	S	3	Sighted	11/08/07	PL	Rusheen, Galway (38 km, SSE, 56 days)
(,	S	3	Sighted	13/10/07	MD	Quilty, Clare (83 km, S, 119 days)
	S	2	Sighted	15/07/09	AOD	Rusheen, Galway (38 km, SSE, 56 days)
			6			, , ,
EW25991	N	1		24/06/07	AB	Rams Island, Mayo
(291V)	S	5	Sighted	16/07/08	DMN	Quilty, Clare (83 km, S, 1 yr 22days)
	S	2	Sighted	13/06/09	EM	Rams Island, Mayo
	S	2	Sighted	26/06/09	EM	Dooghmakeown, SW Mayo